

Is methamphetamine dangerous?

Yes. Methamphetamine causes the heart to beat faster and blood pressure to rise. Since the content of the drug sold varies widely, it is difficult to judge the size of dose. An overdose of methamphetamine can result in seizures, high body temperature, irregular heartbeat, heart attack, stroke and death. The risk of overdose is highest when the drug is injected.

Injecting methamphetamine also puts the user at risk of infections from used needles or impurities in the drug, and of hepatitis or HIV if they share needles with others.

Driving or operating machinery while under the influence of methamphetamine, or any drug, increases the risk of physical injury to the user, and increases the risk of injury to others.

What are the long-term effects of using methamphetamine?

When methamphetamine is used regularly over a long period of time, people can develop amphetamine psychosis. The symptoms of amphetamine psychosis include hallucinations, delusions, paranoia and bizarre and violent behaviour.

Research in animals and humans suggests that methamphetamine may cause long-term damage to cells in those areas of the brain associated with thinking, memory and movement. Further research is needed to determine if these effects are permanent.

One in a series...

Alcohol	Hallucinogens
Alcohol, Other Drugs and Driving	Heroin
Amphetamines	Inhalants
Anabolic Steroids	Ketamine
Benzodiazepines	LSD
Caffeine	Methadone
Cannabis	Methamphetamine
Cocaine	Opioids
Ecstasy	Rohypnol
GHB	Tobacco

For more information on addiction and mental health issues, or a copy of this brochure, please contact CAMH's R. Samuel McLaughlin Addiction and Mental Health Information Centre:
ONTARIO TOLL-FREE: 1 800 463-6273
TORONTO: 416 595-6111

To order multiple copies of this brochure, or other CAMH resource materials, please contact:
Marketing and Sales Services
TEL.: 1 800 661-1111 or 416 595-6059 in Toronto
E-MAIL: marketing@camh.net

To make a donation, please contact:
Centre for Addiction and Mental Health Foundation
TEL.: 416 979-6909
E-MAIL: foundation@camh.net

If you have questions, compliments or concerns about services at CAMH, please call our Client Relations Co-ordinator at:
TEL.: 416 535-8501 ext. 2028

Visit our Web site at: www.camh.net

Copyright © 2003 Centre for Addiction and Mental Health



A Pan American Health Organization / World Health Organization
Collaborating Centre
Affiliated with the University of Toronto

Do You Know...

Street Name: speed, meth, chalk, ice, crystal, crystal meth, jib

What is it?

Methamphetamine belongs to a family of drugs called amphetamines — powerful stimulants that speed up the body's central nervous system. In the 1930s methamphetamine was marketed as a nasal decongestant, and is still medically available in the U.S. as a treatment for obesity. The medical usefulness of methamphetamine is limited by the severity of its adverse effects, and by its high addictive potential. Methamphetamine is not legally available in Canada.



Where does methamphetamine come from?

The methamphetamine that is produced for recreational use is made in illicit labs with fairly inexpensive, and often toxic or flammable, ingredients. The chemicals and processes used vary from lab to lab, affecting the strength, purity and effect of the final product.

What does methamphetamine look like?

Methamphetamine is a white, odourless, bitter-tasting crystalline powder that dissolves easily in water or alcohol and may be snorted, swallowed, smoked or injected. In its smokable form, methamphetamine is called “ice,” “crystal,” “crank” or “glass” because of its transparent, sheet-like crystals. It is smoked in a pipe like crack cocaine.

Who uses methamphetamine?

In the past, illicit methamphetamine use was most closely associated with biker gangs, and also had a spell of popularity in the hippie culture of the 1960s. More recently, the low cost, ease of manufacture and availability of methamphetamine has led to a rise in use among a variety of people. These users include young people at raves, nightclubs and parties, and cocaine users who substitute methamphetamine for its cocaine-like effects.

How does methamphetamine make you feel?

The way methamphetamine — or any other drug — affects you depends on many factors, including:

- . your age and your body weight
- . how much you take and how often you take it
- . how long you’ve been taking it
- . the method you use to take the drug
- . the environment you’re in
- . whether or not you have certain pre-existing medical or psychiatric conditions
- . if you’ve taken any alcohol or other drugs (illicit, prescription, over-the-counter or herbal).

Immediately after smoking methamphetamine or injecting it into a vein, the user experiences an intense surge of euphoria, called a “rush” or “flash.” Snorting methamphetamine produces effects within three to five minutes; swallowing in about 15–20 minutes.

Methamphetamine makes people feel alert and energetic, confident and talkative. They feel little need for food or sleep. On the other hand, users are also likely to feel the many unwanted effects of the drug, including racing of the heart, chest pain, dryness of the mouth, nausea, vomiting and diarrhea and physical tension. Many report an anxious “wired” feeling of restlessness and irritability. The negative effects of methamphetamine can be extreme and alarming, including paranoid delusions, hallucinations, aggressive behaviour and impulsive violence.

How long does the feeling last?

When methamphetamine is injected or taken by mouth, the effects of the drug last about six to eight hours. Smoking methamphetamine may produce effects that last from 10–12 hours. After the effects of the drug have worn off, users are left feeling tired and depressed. Some use the drug continuously over a period of days or weeks in a “binge and crash” pattern, inviting serious health risks, and leading to drug dependency.

Is methamphetamine addictive?

Yes. Tolerance to the effects of methamphetamine builds up quickly in regular users, meaning they need more and more of the drug to achieve the desired effect. When dependent users stop taking methamphetamine, they have strong cravings for the drug, and within a few days will experience withdrawal symptoms, including stomach pain, hunger, headaches, shortness of breath, tiredness and depression.